

**RESTRICTED USE PESTICIDE**

**Toxic to fish and aquatic organisms.**

**For retail sale to and use only by certified applicators, or persons under their direct supervision and only for the uses covered by the certified applicator's certification.**

THIS RECOMMENDATION IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN APPROVED BY OR SUBMITTED TO EPA FOR APPROVAL.



## 2(ee) Recommendation

**FOR COMMERCIAL NON-FOOD USE IN INTERIORSCAPES AND ON OUTDOOR ORNAMENTALS, CHRISTMAS TREES, & NURSERIES TO CONTROL SPOTTED LANTERNFLY**

**THIS 2(ee) RECOMMENDATION IS FOR DISTRIBUTION AND USE IN THE STATE OF NEW YORK.**

**IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.**

This product bulletin contains new or additional directions for use recommended by FMC Corp. under FIFRA Section 2(ee).

<b>Ornamental and Tree Foliar Application Rates</b>	<b>PEST</b>	<b>RATE OF APPLICATION</b>
Ornamentals in Interiorscapes and on Outdoor Ornamentals, Christmas Trees, & Nurseries	Spotted Lanternfly	0.08 – 0.16 fl oz/1000 sq ft or 3.6 – 7.2 fl oz/100 gal (0.056 – 0.11 lb ai/100 gal)

### RESTRICTIONS AND PRECAUTIONS

At the discretion of the applicator, this product may be applied at up to 0.32 fluid oz. per 1,000 square feet (14.4 fl. oz. per 100 gallons) to control spotted lanternfly. The higher application rates should be used when maximum residual control is desired.

Certain cultivars may be sensitive to the final spray solution. A small number of plants should be treated and observed for one week prior to application to the entire planting.

**In New York State, this product may NOT be applied to any grass or turf area within 100 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch).**

**In New York State, do make a single repeat application of this product if there are signs of renewed insect activity, but not sooner than two weeks after the first application.**

- Do not apply to pets, crops, or sources of electricity.
- Firewood is not to be treated.
- Do not allow spray to contact food, foodstuffs, food contacting surfaces, food utensils or water supplies.
- Do not apply this pesticide in livestock buildings (barns).
- Keep children and pets off treated areas following application until the spray has dried.
- Do not apply by air.
- Do not use in greenhouses.
- Do not apply this product through any type of irrigation system.
- Do not apply when a temperature inversion exists.
- Do not apply for surface feeding pests if rain is expected within 12 hours (or whatever time is necessary for the spray to dry).
- Do not apply within 25 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish farm ponds.
- Do not apply when grass areas are water logged or the soil is saturated with water (i.e. will not accept irrigation).
- Do not use on golf courses or sod farms on in Nassau or Suffolk Counties, NY

- **Vinyl and Aluminum Siding:** Do not spray directly onto vinyl or aluminum siding. If OnyxPro inadvertently contacts vinyl and aluminum siding (particularly lightly colored, aged, weathered or otherwise damaged), it may result in staining, bleaching or discoloration. Wash off thoroughly with detergent and water. Factors such as extreme heat and direct sunlight can promote damage when using emulsifiable concentrates. Avoid application to vinyl or aluminum siding while exposed to direct sunlight or during the heat of the day.

**ALL APPLICABLE DIRECTIONS, RESTRICTIONS, WORKER PROTECTION STANDARDS REQUIREMENTS AND PRECAUTIONS ON THE EPA-REGISTERED ONYXPRO, EPA REG. NO. 279-4269, LABEL MUST BE FOLLOWED. THE ONYXPRO SECTION 3 LABEL AND THIS RECOMMENDATION MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.**



FMC Corporation  
Agricultural Products Group  
2929 Walnut Street  
Philadelphia PA 19104

2/2019