



Onyx Insecticide

For use in control of *Dendroctonus* bark beetles in trees grown for non-commercial purposes.

EPA REG No. 279-3177

EPA SLN No. SD- 130002

24(c) Supplemental Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF SOUTH DAKOTA

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This label must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

This 24(c) label is only approved for use in State of South Dakota on trees grown for non-commercial purposes, including trees on private lands, parks, or rangeland. Not for use on trees being grown for sale or other commercial use, for commercial seed production, or trees grown for research purposes.

TRUNK SPRAYS

For control of *Dendroctonus* bark beetles such as mountain pine beetle, southern pine beetle, western pine beetle, Black turpentine beetle, and engraver beetles (*Ips* spp.)

Preventative control: Make applications of a spray mixture containing 1.0 to 2.0 pints of this product per 100 gallons (0.25 to 0.5 lbs. ai/100 gallons) of water to the trunk of the tree with a hydraulic sprayer in the spring of the year or when a threat of infestation is evident from nearby infested trees.

Apply spray directly to the main trunk from the base of the tree to at least half way into the live crown. Spray until the bark is thoroughly wetted by the spray (usually 1 to 4 gallons of spray per tree).

Application rates and application timing differ according to the target pest and other factors peculiar to each local situation. Consult your local State Extension specialist or other qualified expert for specific recommendations.

Treatment of Infested trees to control emerging brood: Make applications of a spray mixture containing 2.0 pints of this product per 100 gallons of water to trees that still have beetles in the bark.

Apply spray directly to the main trunk from the base of the tree to at least half way into the live crown. Spray until the bark is thoroughly wetted by the spray (usually 1 to 4 gallons of spray per tree).

Trees on which all needles have turned brown generally have been vacated and should not be sprayed unless infestation is confirmed. To confirm an infestation, scrape off the outer bark to determine if trees are still infested. If live infestations remain in the trunks, fell the trees and cut into sections. Spray the trunk and large limbs and turn sections so that all of the surface area can be treated.

Restrictions: Do not apply more than 0.6 lbs active ingredient per acre

For MEDICAL Emergencies only, have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-(800)-331-3148 for Emergency Assistance.

For Information Regarding the Use of this Product Call 1-800-321-1FMC (1362).
As with any pesticide, always read and follow label directions.

Onyx is a registered trademark of FMC Corporation

FMC Corporation
Agricultural Products Group
1735 Market Street
Philadelphia, PA 19103