

**RESTRICTED USE PESTICIDE**

**Toxic to fish and aquatic organisms.**

**For retail sale to and use only by certified applicators, or persons under their direct supervision and only for the uses covered by the certified applicator's certification.**

THIS RECOMMENDATION IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN APPROVED BY OR SUBMITTED TO EPA FOR APPROVAL.



## 2(ee) Recommendation

**FOR COMMERCIAL NON-FOOD USE ON INDOOR AND OUTDOOR ORNAMENTALS IN GREENHOUSES AND NURSERIES TO CONTROL SPOTTED LANTERNFLY**

**THIS 2(ee) RECOMMENDATION IS FOR DISTRIBUTION AND USE IN THE STATE OF NEW YORK.**

**IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.**

This product bulletin contains new or additional directions for use are recommended by FMC Corp. under FIFRA Section 2(ee).

<b>Ornamental Application Directions</b>	<b>PEST</b>	<b>RATE OF APPLICATION</b>
Ornamentals in Greenhouses, Lath Houses, Shade Houses and Outdoor Nurseries, including Non-Bearing Fruit and Nut Trees	Spotted Lanternfly	10 - 20 Fl oz/A (0.05 to 0.10 Lb ai/A)

### RESTRICTIONS AND PRECAUTIONS

Talstar Select Insecticide may be diluted and applied in various volumes of water providing that the maximum label rate (0.2 lbs ai/A or 40 fl. ozs.) is not exceeded (refer to Dilution Chart on the full label for specific instructions). Talstar Select Insecticide may be applied through low volume application equipment by dilution with water or other carriers and providing that the maximum label rate (0.2 lbs ai/A or 40 fl. ozs.) is not exceeded.

Certain cultivars may be sensitive to the final spray solution. A small number of plants should be treated and observed for one week prior to application to the entire planting.

**In New York State, this product may NOT be applied to any grass or turf area within 100 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch).**

**In New York State, do make a single repeat application of Talstar Select Insecticide if there are signs of renewed insect activity, but not sooner than two weeks after the first application.**

- Apply with ground equipment only.
- Do not apply when wind direction favors downwind drift towards nearby water bodies.
- Do not apply when wind velocity exceeds 10 mph.
- Avoid application when wind gusts approach 10 mph.
- Do not apply when a temperature inversion exists.
- Apply using the largest nozzle size compatible with adequate coverage.
- Do not apply if rain is expected within 12 hours (or whatever time is necessary for the spray to dry).

- Do not apply by ground equipment within 25 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries, and commercial fish farm ponds. When treating tall trees (>15 feet) from the ground with high pressure sprays or during any application with air assisted equipment (mist blower) do not apply within 150 feet of aquatic areas.

**ALL APPLICABLE DIRECTIONS, RESTRICTIONS, WORKER PROTECTION STANDARDS REQUIREMENTS AND PRECAUTIONS ON THE EPA-REGISTERED TALSTAR S, EPA REG. NO. 279-3155, LABEL MUST BE FOLLOWED. THE TALSTAR S SECTION 3 LABEL AND THIS RECOMMENDATION MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.**



FMC Corporation  
Agricultural Products Group  
2929 Walnut Street  
Philadelphia PA 19104

2/2019