

Supplemental Labeling



EPA Reg. No. 8033-96-279

Active Ingredient:	By Wt.
Acetamiprid.....	22.73 %
Bifenthrin*	27.27 %
Inert Ingredients:	50.00 %
	100.00%

*Cis isomers 97% minimum, trans isomers 3% maximum.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL ARE TO BE FOLLOWED.

This Supplemental labeling must be in the possession of the user at the time of pesticide application. Read the label affixed to the container for Transport Termiticide Insecticide before applying. Carefully follow all precautionary statements and application use directions. Use of Transport Termiticide Insecticide according to this supplemental labeling is subject to all use precautions and limitations imposed by the labeling affixed to the container for Transport Termiticide Insecticide.

FMC

FMC Corporation
Agricultural Products Group
1735 Market Street
Philadelphia PA 19103

This product can also be used to control ants and a range of other general household pests outdoors around the exterior perimeter of buildings and structures.

STORAGE AND DISPOSAL

Pesticide Storage: Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Do not store at temperatures below 32°F (0°C). Rough handling may cause breakage, especially at low temperatures. Allow warming above 50°F (10°C) before use. Do not allow inner bags to become wet during storage. Do not handle inner bag with hands or wet gloves. Do not put concentrate or diluted material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food, or feed by storage or disposal.

In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills.

To Confine Spill: If liquid, dike surrounding area or absorb with sand, cat litter or commercial clay. If dry material, cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

Pesticide Disposal: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. Dispose of excess or waste pesticide by use according to label directions, or contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal:

Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available or dispose of empty outer pouch in the trash as long as water soluble bags were intact.

APPLICATION RATE CHART

IF USING 7.5 OUNCE BAG

Number of Gallons Desired	TERMITE USES		GENERAL HOUSEHOLD PEST USES	
	Number of 7.5 ounce bags	Dilution Rate (% Active Ingredient)	Number of 7.5 ounce bags	Dilution Rate (% Active Ingredient)
25	1	0.11%	1	0.11%
50	2	0.11%	2	0.11%
75	3	0.11%	3	0.11%
100	4	0.11%		

Dilution Chart

Dilution Chart for Perimeter Barrier Applications Around Structures

Application Volume per 1,000 sq. ft.	Transport 7.5 ounce water soluble bags to add			
	Total Mix volume			
	25 Gallons	50 Gallons	75 Gallons	100 Gallons
1.0 gal/1,000 sq. ft	1 packet	2 packets	3 packets	4 packets

Foam Applications

The Transport Termiticide Insecticide dilution may be converted to foam with expansion characteristics from 2 to 40 times for localized control or prevention of pests including ants, bees, termites, wasps or other arthropods harboring in walls, under slabs or in other void areas.

Depending on the circumstances, foam applications may be used alone or in combination with liquid dilution applications. Applications may be made behind veneers, piers, chimney bases, into rubble foundations, into block voids or structural voids, under slabs, stoops, porches, or to the soil in crawlspaces, and other similar voids.

Foam and liquid application must be consistent with volume and active ingredient instructions in order to insure proper application has been made. The volume and amount of active ingredient are essential to an effective treatment. At least 75% of the labeled liquid dilution volume of product must be applied, with the remaining percent delivered to appropriate areas using foam application. Refer to label and use recommendations of the foam manufacturer and the foaming equipment manufacturer.

Foam applications are generally a good supplement to liquid treatments in difficult areas, but may be used alone in difficult spots.

Use dry foam (20:1 or greater expansion ratio) for applications to wall voids and stud walls.

Use wet foam (5:1 or lower expansion ratio) for applications to soil, including applications to filled porches.

Application Under Slabs or to Soil in Crawlspaces to Prevent or Control Termites and other pests	Application may be made using Transport Termiticide Insecticide foam alone or in combination with liquid dilution. The equivalent of at least 4 gallons of dilution per 10 linear feet (vertical barrier), or at least 1 gallon of dilution per 10 square feet (horizontal barrier) must be applied either as dilution, foam, or a combination of both.
---	---

General Household Pest Control – Outdoor Pinstream, Spot, Crack and Crevice or Perimeter Treatment

Controls	Ants (including Red Imported Fire Ants and Carpenter Ants) Bees Beetles* Biting Flies Boxelder Bugs Centipedes Chiggers Clover Mites Cockroaches Crickets Earwigs Elm Leaf Beetles Firebrats Fleas Flies	Gnats Midges Millipedes Mosquitoes Moths Pillbugs Scorpions Silverfish Sowbugs Spider Mites Spiders (including Black Widow) Springtails Ticks (including American Dog Ticks and Deer Ticks) Wasps *Not for use in California
Where to Apply	Apply as a pinstream, spot, crack and crevice, or perimeter spray on and around outside surfaces of residential and non-residential buildings and structures including, but not limited to, exterior siding, foundations, porches, window frames, eaves, patios, garages, areas adjacent or around private homes, duplexes, townhouses, condominiums, house trailers, apartment complexes, carports, garages, storage sheds, and other structures, and other areas where pests congregate or have been seen.	
Perimeter Treatment Application Rate	As a perimeter treatment, apply as a continual band up to 10 foot wide around the structure and upwards along the foundation to a height of up to 3 feet and around windows, doors, other penetrations and roof eaves, soffets and overhangs. Apply 0.11% concentration of Transport Termiticide Insecticide in sufficient amount of water (see Dilution Chart) and apply adequately to cover 1,000 square feet. Dilutions may be applied at either high or low volumes. Do not apply more than 0.11% finished dilution per 1,000 square feet.	
Mixing Directions	Fill tank 1/4 to 1/3 full with water. Start pump to begin by-pass agitation and place end of treating tool in tank to allow circulation through hose. Add Transport Termiticide Insecticide bag(s). Add remaining amount of water. Let pump run and allow recirculation through the hose for 2 to 3 minutes. Transport Termiticide Insecticide may also be mixed into full tanks of water, but requires substantial agitation to insure uniformity of the dilution.	
Repeat Applications	Retreatment may be necessary to achieve and/or maintain control during periods of high pest pressure. Repeat application is necessary only if there are signs of renewed insect activity.	

Outdoor Ant Control

Carpenter Ants	Apply Transport Termiticide Insecticide as a pinstream, spot, crack and crevice, or perimeter spray to carpenter ant trails around doors and windows and other places where carpenter ants have been observed or are expected to forage. For best results, locate and treat carpenter ant nests. Apply a perimeter treatment using either low or high volume applications described in the General Household Pest Control - Outdoor section of this label. The higher dilutions and/or application volumes, as well as more frequent applications, may be necessary when treating concrete surfaces for ant control. Maximum control is generally achieved using the following procedure: 1) Treat non-porous surfaces with low volume applications using 0.11% dilution per 1,000 square feet. 2) Treat porous surfaces and vegetation with high volume applications using 0.11% dilution per 1,000 square feet. 3) Treat the trunks of trees that have carpenter ant trails or upon which carpenter ants are foraging using 0.11% dilution and applying this dilution to thoroughly wet the bark from the base of the tree to as high as possible on the trunk.
Nuisance Ants Outdoor (including fire ants)	For best results, locate and treat ant nests. Apply Transport Termiticide Insecticide as a pinstream, spot, crack and crevice or perimeter treatment to ant trails around doors and windows and other places where ants have been observed or are expected to forage. Apply a perimeter treatment described in the General Household Pest Control - Outdoor section of this label. More frequent applications, may be necessary when treating concrete surfaces for ant control. Maximum control is generally achieved using the following procedure: 1) Treat non-porous surfaces with low volume applications using 0.11% dilution per 1,000 square feet. 2) Treat porous surfaces and vegetation with high volume applications using 0.11% dilution per 1,000 square feet.

Specific Pest Control Applications

Ant and Fire Ant Mounds	Drench mounds with 1-2 gallons of Transport Termiticide Insecticide at a 0.11% dilution (see Dilution Chart) to each mound area by sprinkling the mound until it is wet and treat 3-feet out around the mound. Use the higher volume for mounds larger than 12". For best results, apply in cool weather, such as in early morning or late evening hours, but not in the heat of the day.
Carpenter Ants in trees, utility poles, fencing, deck materials and similar structural members	Drill to locate the interior infested cavity and inject or foam the 0.11% dilution (see Dilution Chart) into the cavity using a sufficient volume and an appropriate treatment tool with a splash-back guard.
Wood piles and stored lumber	To protect firewood or lumber from carpenter ants, make up a 0.11% dilution (see Dilution Chart) of Transport Termiticide Insecticide and apply as a spot treatment to the soil beneath where the firewood or lumber will be stacked at the rate of one gallon of dilution per 8 square feet. Treated wood can be burned as firewood or used as lumber one month after treatment.
Underground Services	Underground Services such as: wires, cables, utility lines, pipes, conduits, etc. Services may be within structures or located outside of structures. Soil treatment may be made using Transport Termiticide Insecticide dilution to prevent attack by Termites and Ants. Apply 2 gallons of 0.11% dilution (see Dilution Chart) per 10 linear feet to the bottom of the trench and allow liquid to soak into the soil. Lay services on the treated soil and cover with approximately 2 inches of fill soil. Apply another 2 gallons per 10 linear feet over the soil surface to complete the treatment barrier. In wide trenches, only treat the soil in the area near the services. It is important to establish a continuous barrier of treated soil surrounding the services. Where soil will not accept the above-labeled volume, 1 gallon of 0.11% dilution of Transport Termiticide Insecticide may be used per 10 linear feet of trench both to the bottom of the trench and over the soil on top of the services. Finish filling the trench with treated fill soil. The soil where each service protrudes from the ground may be treated by trenching/rodding of no more than 1 to 2 gallons of 0.11% dilution into the soil.
Posts, Poles and Other Constructions	Create an insecticidal barrier in the soil around wooden constructions such as signs, fences and landscape ornamentation by applying a dilution. Previously installed poles and posts may be treated by sub-surface injection or treated by gravity-flow through holes made from the bottom of a trench around the pole or post. Treat on all sides to create a continuous insecticidal barrier around the pole. Use 1 gallon of 0.11% dilution (see Dilution Chart) per foot of depth for poles and posts less than six inches in diameter. For larger poles, use 1.5 gallons of 0.11% dilution per foot of depth. Apply to a depth of 6 inches below the bottom of the wood. For larger constructions, use 4 gallons per 10 linear feet per foot of depth.
Pests Under Slabs	Infestations of Arthropods, such as Ants, Cockroaches and Scorpions inhabiting under slab areas may be controlled by drilling and injecting or horizontal rodding and then injecting 1 gallon of 0.11% dilution (see Dilution Chart) per 10 square feet or 2 gallons of 0.11% dilution per 10 linear feet.
Pest Control in Crawlspace and Voids	Apply Transport Termiticide/Insecticide 0.11% dilution (see Dilution Chart) to all surfaces in crawlspace and/or voids to control ants, fleas, roaches, scorpions, or other arthropods. Product may also be applied through insecticidal delivery systems such as piping or flexible tubing mounted under and/or around the structure as a crack and crevice or spot treatment. Treat surfaces to point of runoff. Keep children and pets off surface until dry.

General Household Pest Control - Indoor

Controls	Ants (including Red Imported Fire Ants and Carpenter Ants) Bed Bugs Bees Beetles (including Carpet Beetles) Boxelder Bugs Centipedes Cockroaches Crickets Earwigs Firebrats Fleas Flies	Gnats Midges Millipedes Moths (including Cloth Moths) Pillbugs Scorpions Silverfish Sowbugs Spiders (including Black Widow and Brown Recluse) Ticks Wasps
-----------------	--	---

Where to Apply	Apply for residual pest control in residential and non-residential buildings and structures. Apply either as a crack and crevice, pinstream, spot, coarse, low-pressure spray (25 p.s.i. or less) or with a paintbrush. Apply to areas where pests hide, such as baseboards, corners, storage areas, closets, around water pipes, doors and windows, attics and eaves, behind and under refrigerators, dishwashers, cabinets, sinks, furnaces, stoves, the underside of shelves, drawers and similar areas and other possible pest harborage sites. Do not use as a space spray. Pay particular attention to cracks and crevices.
Application Rate	Apply 0.11% concentration of active ingredient in sufficient amount of water (see Dilution Chart) and apply adequately to cover 1,000 square feet. Do not apply more than 0.11% finished dilution per 1,000 square feet.
Mixing Directions	Fill tank 1/4 to 1/3 full with water. Start pump to begin by-pass agitation and place end of treating tool in tank to allow circulation through hose. Add Transport Termiticide Insecticide bag(s). Add remaining amount of water. Let pump run and allow recirculation through the hose for 2 to 3 minutes. Transport Termiticide Insecticide may also be mixed into full tanks of water, but requires substantial agitation to insure uniformity of the dilution.
Ants (including Carpenter Ants and Nuisance Ants)	Apply to areas where ants have been observed or are expected to forage. Particular attention should be given to treating entry points into the home or premises such as around doors and windows and other places where ants and ant trails may be found. For added Carpenter Ant control, spray or foam into cracks and crevices or drill holes and spray, mist or foam into voids where Carpenter Ants or their nests are present. When using Transport Termiticide Insecticide in combination with baits, apply Transport Termiticide Insecticide as instructed above, and use baits in other areas that have not been treated with Transport Termiticide Insecticide.
Bed Bugs	Thorough application should be made to crack and crevices where evidence of bed bugs occurs. This includes bed frames, box springs, inside empty dressers and clothes closets and carpet edges, high and low wall moldings and wallpaper edges. Do not use this product on bed linens, pillows, mattresses or clothes. Remove all clothes and other articles from dressers or clothes closets before application. Allow all treated areas to thoroughly dry before use. Not recommended for use as sole protection against bedbugs. If evidence of bedbugs is found in/on mattresses, use products approved for this use.
Bees and Wasps	To control Bees, Wasps, Hornets, and Yellow Jackets indoors, application should be made in the late evening when insects are at rest. Spray liberally into hiding and breeding places, especially under attic rafters, contacting as many insects as possible.
Occasional Invaders	To control Boxelder Bugs, Centipedes, Earwigs, Beetles, Millipedes, Pillbugs, and Sowbugs, apply around doors and windows and other places where these pests may be found or where they may enter premises. Spray baseboards, storage areas and other locations.
Crawling and Flying Insect Pests	To control Cockroaches, Crickets, Firebrats, Flies, Gnats, Midges, Moths, Scorpions, Silverfish, Spiders, apply as a coarse, low pressure spray to areas where these pests hide, such as baseboards, corners, storage areas, closets, around water pipes, doors and windows, attics and eaves, behind and under dishwashers, refrigerators, cabinets, sinks, furnaces, and stoves, the underside of shelves, drawers and similar areas. Pay particular attention to cracks and crevices.
Important	Do not use in food/feed areas of food/feed handling establishments, restaurants, or other areas where food is commercially prepared or processed. Do not use in serving areas while food/feed is exposed or facility is in operation. Serving areas are areas where prepared foods are served, such as dining rooms, but excluding areas where food may be prepared or held. In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Exposed food should be covered or removed. Not for use in Federally inspected meat and poultry plants.

Specific Indoor Pest Control Applications

Warehouses and Stores	Transport Termiticide Insecticide may be applied as a spot or crack and crevice treatment in non-food storage warehouses and stores. Apply no more than 0.11% dilution (see Dilution Chart) of Transport Termiticide Insecticide per 1,000 square feet to provide adequate coverage. Apply to all areas that may harbor pests, including under and between pallets, bins and shelves. Do not apply directly to food grain bins (interior) or animals.
------------------------------	---

RESTRICTIONS

Do not apply in a perimeter treatment to areas beyond 10 feet from the foundation of the structure.

Do not use as a space spray.

Do not use in and around the exterior perimeter of commercial barns, stables, paddocks. Do not use in grazing areas, feed lots or other similar areas used for housing, boarding, and/or rearing animals.

Do not apply by air.

Do not apply as a broadcast spray on lawns and turf.

Do not apply in greenhouses or nurseries.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.

Do not apply this product through any kind of irrigation system.

Not for use on sod farm turf, golf course turf, or grass grown for seed.

Do not apply to pets, crops, or sources of electricity.

Do not treat electrically active underground services.

Do not treat areas where food is exposed. Do not allow spray to contact food, foodstuffs, food contacting surfaces, food utensils or water supplies.

Thoroughly wash dishes and food handling utensils with soap and water if they become contaminated by application of this product.

Let surfaces dry before allowing people and pets to contact surfaces.

Transport Termiticide Insecticide will not stain or damage any surface that water alone will not stain or damage.

Application equipment that delivers low volume treatments, such as the Micro-Injector® or Actisol® applicators, may also be used to make crack and crevice, deep harborage, spot and general surface treatments of Transport Termiticide Insecticide.

Wear protective clothing; unvented goggles, gloves and a respirator approved by NIOSH, when applying to overhead areas or in poorly ventilated or confined areas.

Conditions of Sale of Warranty and Liability:

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions beyond the control of FMC Corporation or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold FMC Corporation and Seller harmless for any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the Directions for Use when used in accordance with the directions under normal conditions of use. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, FMC Corporation MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE, NOR ANY OTHER EXPRESS OR IMPLIED WARRANTIES WITH RESPECT TO THE SELECTION, PURCHASE, OR USE OF THIS PRODUCT TO THE EXTENT ALLOWED BY STATE LAW. Any warranties, express or implied, having been made are inapplicable if this product has been used contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to (or beyond the control of) seller or FMC Corporation and buyer assumes the risk of any such use.

To the extent consistent with applicable law, FMC Corporation or seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF FMC Corporation AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF FMC Corporation OR SELLER, THE REPLACEMENT OF THE PRODUCT.

This Conditions of Sale and Limitation of Warranty and Liability may not be amended by any oral or written agreement.

FMC - Trademark of FMC Corporation

Transport - Trademark of Nippon Soda Co., Ltd